NDA & NA Exam (1), 2021

जब तक आपको यह परीक्षण पुस्तिका खोलने को न कहा जाए तब तक न खोलें

टी.बी.सी. : FJY-D-MTC

परीक्षण पुस्तिका अनुक्रम

क्रम संख्या

1333833

परीक्षण पुस्तिका गणित



समय : दो घण्टे और तीस मिनट

पूर्णांक : 300

अ नु दे श

- परीक्षा प्रारम्भ होने के तुरन्त बाद आप इस परीक्षण पुस्तिका की पड़ताल अवश्य कर लें कि इसमें कोई बिना छपा, फटा या छूटा हुआ पृष्ठ अथवा प्रश्नांश आदि न हो। यदि ऐसा है, तो इसे सही परीक्षण पुस्तिका से बदल लें।
- 2. कृपया ध्यान रखें कि OMR उत्तर-पत्रक में उचित स्थान पर रोल नम्बर और परीक्षण पुस्तिका अनुक्रम A, B, C या D को ध्यान से एवं बिना किसी चूक या विसंगति के भरने और कूटबद्ध करने की जिम्मेदारी उम्मीदवार की है। किसी भी प्रकार की चूक/विसंगति की स्थिति में उत्तर-पत्रक निरस्त कर दिया जाएगा।
- इस परीक्षण पुस्तिका पर साथ में दिए गए कोष्ठक में आपको अपना अनुक्रमांक लिखना है। परीक्षण पुस्तिका पर और कुछ न लिखें।
- 4. इस परीक्षण पुस्तिका में 120 प्रश्नांश (प्रश्न) दिए गए हैं। प्रत्येक प्रश्नांश हिन्दी और अंग्रेजी दोनों में छपा है। प्रत्येक प्रश्नांश में चार प्रत्युत्तर (उत्तर) दिए गए हैं। इनमें से एक प्रत्युत्तर को चुन लें, जिसे आप उत्तर-पत्रक पर अंकित करना चाहते हैं। यदि आपको ऐसा लगे कि एक से अधिक प्रत्युत्तर सही हैं, तो उस प्रत्युत्तर को अंकित करें जो आपको सर्वोत्तम लगे। प्रत्येक प्रश्नांश के लिए केवल एक ही प्रत्युत्तर चुनना है।
- 5. आपको अपने सभी प्रत्युत्तर अलग से दिए गए उत्तर-पत्रक पर ही अंकित करने हैं। उत्तर-पत्रक में दिए गए निर्देश देखें।
- 6. सभी प्रश्नांशों के अंक समान हैं।
- 7. इससे पहले कि आप परीक्षण पुस्तिका के विभिन्न प्रश्नांशों के प्रत्युत्तर उत्तर-पत्रक पर अंकित करना शुरू करें, आपको प्रवेश प्रमाण-पत्र के साथ प्रेषित अनुदेशों के अनुसार कुछ विवरण उत्तर-पत्रक में देने हैं।
- 8. आप अपने सभी प्रत्युत्तरों को उत्तर-पत्रक में भरने के बाद तथा परीक्षा के समापन पर केवल उत्तर-पत्रक अधीक्षक को सौंप दें। आपको अपने साथ परीक्षण पुस्तिका ले जाने की अनुमति है।
- 9. कच्चे काम के लिए पत्रक, परीक्षण पुस्तिका के अन्त में संलग्न हैं।
- 10. गलत उत्तरों के लिए दण्ड :

वस्तुनिष्ठ प्रश्न-पत्रों में उम्मीदवार द्वारा दिए गए गलत उत्तरों के लिए दण्ड दिया जाएगा।

- (i) प्रत्येक प्रश्न के लिए चार वैकल्पिक उत्तर हैं। उम्मीदवार द्वारा प्रत्येक प्रश्न के लिए दिए गए एक गलत उत्तर के लिए प्रश्न हेतु नियत किए गए अंकों का एक-तिहाई दण्ड के रूप में काटा जाएगा।
- (ii) यदि कोई उम्मीदवार एक से अधिक उत्तर देता है, तो इसे गलत उत्तर माना जाएगा, यद्यपि दिए गए उत्तरों में से एक उत्तर सही होता है, फिर भी उस प्रश्न के लिए उपर्युक्तानुसार ही उसी तरह का दण्ड दिया जाएगा।
- (iii) यदि उम्मीदवार द्वारा कोई प्रश्न हल नहीं किया जाता है अर्थात् उम्मीदवार द्वारा उत्तर नहीं दिया जाता है, तो उस प्रश्न के लिए कोई दण्ड नहीं दिया जाएगा।

जब तक आपको यह परीक्षण पुस्तिका खोलने को न कहा जाए तब तक न खोलें

1. The smallest positive integer n for which

$$\left(\frac{1-i}{1+i}\right)^{n^2} = 1$$

where $i = \sqrt{-1}$, is

- (a) 2
- (b) 4
- (c) 6
- (d) 8
- 2. The value of x, satisfying the equation $\log_{\cos x} \sin x = 1$, where $0 < x < \frac{\pi}{2}$, is
 - (a) $\frac{\pi}{12}$
 - (b) $\frac{\pi}{3}$
 - (c) $\frac{\pi}{4}$
 - (d) $\frac{\pi}{6}$
- 3. If Δ is the value of the determinant

$$\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

then what is the value of the following determinant?

$$\begin{vmatrix} pa_1 & b_1 & qc_1 \\ pa_2 & b_2 & qc_2 \\ pa_3 & b_3 & qc_3 \end{vmatrix}$$

 $(p \neq 0 \text{ or } 1, q \neq 0 \text{ or } 1)$

- (a) $p\Delta$
- (b) q∆
- (c) $(p+q)\Delta$
- (d) $pq\Delta$

- **4.** If C_0 , C_1 , C_2 , ..., C_n are the coefficients in the expansion of $(1+x)^n$, then what is the value of $C_1 + C_2 + C_3 + \cdots + C_n$?
 - (a) 2^n
 - (b) $2^n 1$
 - (c) 2^{n-1}
 - (d) $2^n 2$
- 5. If a+b+c=4 and ab+bc+ca=0, then what is the value of the following determinant?

- (a) 32
- (b) -64
- (c) -128
- (d) 64
- 6. The number of integer values of k, for which the equation $2\sin x = 2k + 1$ has a solution, is
 - (a) zero
 - (b) one
 - (c) two
 - (d) four

7. If $a_1, a_2, a_3, \dots, a_9$ are in GP, then what is the value of the following determinant?

- (a) 0
- (b) 1
- (c) 2
- (d) 4
- **8.** If the roots of the quadratic equation $x^2 + 2x + k = 0$ are real, then
 - (a) k < 0
 - (b) k≤0
 - (c) k < 1
 - (d) $k \le 1$
- **9.** If n = 100!, then what is the value of the following?

$$\frac{1}{\log_2 n} + \frac{1}{\log_3 n} + \frac{1}{\log_4 n} + \dots + \frac{1}{\log_{100} n}$$

- (a) 0
- (b) 1
- (c) 2
- (d) 3

- **10.** If Z = 1 + i, where $i = \sqrt{-1}$, then what is the modulus of $Z + \frac{2}{Z}$?
 - (a) 1
 - (b) 2
 - (c) 3
 - (d) 4
- 11. If A and B are two matrices such that AB is of order $n \times n$, then which one of the following is correct?
 - (a) A and B should be square matrices of same order.
 - (b) Either A or B should be a square matrix.
 - (c) Both A and B should be of same order.
 - (d) Orders of A and B need not be the same.
- 12. How many matrices of different orders are possible with elements comprising all prime numbers less than 30?
 - (a) 2
 - (b) 3
 - (c) 4
 - (d) 6

13. Let

$$A = \begin{vmatrix} p & q \\ r & s \end{vmatrix}$$

where p, q, r and s are any four different prime numbers less than 20. What is the maximum value of the determinant?

- (a) 215
- (b) 311
- (c) 317
- (d) 323
- 14. If A and B are square matrices of order 2 such that det(AB) = det(BA), then which one of the following is correct?
 - (a) A must be a unit matrix.
 - (b) B must be a unit matrix.
 - (c) Both A and B must be unit matrices.
 - (d) A and B need not be unit matrices.

15. What is

 $\cot 2x \cot 4x - \cot 4x \cot 6x - \cot 6x \cot 2x$

equal to?

- (a) -1
- (b) 0
- (c) 1
- (d) 2

- **16.** If $\tan x = -\frac{3}{4}$ and x is in the second quadrant, then what is the value of $\sin x \cdot \cos x$?
 - (a) $\frac{6}{25}$
 - (b) $\frac{12}{25}$
 - (c) $-\frac{6}{25}$
 - (d) $-\frac{12}{25}$
- 17. What is the value of the following?

$$\csc\left(\frac{7\pi}{6}\right)\sec\left(\frac{5\pi}{3}\right)$$

- (a) $\frac{4}{3}$
- (b) 4
- (c) -4
- (d) $-\frac{4}{\sqrt{3}}$
- 18. If the determinant

$$\begin{vmatrix} x & 1 & 3 \\ 0 & 0 & 1 \\ 1 & x & 4 \end{vmatrix} = 0$$

then what is x equal to?

- (a) -2 or 2
- (b) -3 or 3
- (c) -1 or 1
- (d) 3 or 4

19. What is the value of the following?

tan 31° tan 33° tan 35° ··· tan 57° tan 59°

- (a) -1
- (b) 0
- (c) 1
- (d) 2
- 20. If

$$f(x) = \begin{vmatrix} 1 & x & x+1 \\ 2x & x(x-1) & x(x+1) \\ 3x(x-1) & 2(x-1)(x-2) & x(x+1)(x-1) \end{vmatrix}$$

then what is f(-1) + f(0) + f(1) equal to?

- (a) 0
- (b) 1
- (c) 100
- (d) -100
- **21.** The equation $\sin^{-1} x \cos^{-1} x = \frac{\pi}{6}$ has
 - (a) no solution
 - (b) unique solution
 - (c) two solutions
 - (d) infinite number of solutions

22. What is the value of the following?

 $(\sin 24^{\circ} + \cos 66^{\circ})(\sin 24^{\circ} - \cos 66^{\circ})$

- (a) -1
- (b) 0
- (c) 1
- (d) 2
- 23. A chord subtends an angle 120° at the centre of a unit circle. What is the length of the chord?
 - (a) $\sqrt{2}-1$ units
 - (b) $\sqrt{3}-1$ units
 - (c) $\sqrt{2}$ units
 - (d) $\sqrt{3}$ units
- 24. What is

 $(1 + \cot \theta - \csc \theta)(1 + \tan \theta + \sec \theta)$

equal to?

- (a) 1
- (b) 2
- (c) 3
- (d) 4

25. What is

$$\frac{1+\tan^2\theta}{1+\cot^2\theta} - \left(\frac{1-\tan\theta}{1-\cot\theta}\right)^2$$

equal to?

- (a) 0
- (b) 1
- (c) 2 tan θ
- (d) 2cotθ
- 26. What is the interior angle of a regular octagon of side length 2 cm?
 - (a) $\frac{\pi}{2}$
 - (b) $\frac{3\pi}{4}$
 - (c) $\frac{3\pi}{5}$
 - (d) $\frac{3\pi}{8}$
- 27. If $7 \sin \theta + 24 \cos \theta = 25$, then what is the value of $(\sin \theta + \cos \theta)$?
 - (a) 1
 - (b) $\frac{26}{25}$
 - (c) $\frac{6}{5}$
 - (d) $\frac{31}{25}$

- 28. A ladder 6 m long reaches a point 6 m below the top of a vertical flagstaff. From the foot of the ladder, the elevation of the top of the flagstaff is 75°. What is the height of the flagstaff?
 - (a) 12 m
 - (b) 9 m
 - (c) $(6+\sqrt{3})$ m
 - (d) $(6+3\sqrt{3})$ m
- **29.** The shadow of a tower is found to be x metre longer, when the angle of elevation of the sun changes from 60° to 45° . If the height of the tower is $5(3+\sqrt{3})$ m, then what is x equal to?
 - (a) 8 m
 - (b) 10 m
 - (c) 12 m
 - (d) 15 m
- 30. If $3\cos\theta = 4\sin\theta$, then what is the value of $\tan(45^\circ + \theta)$?
 - (a) 10
 - (b) 7
 - (c) $\frac{7}{2}$
 - (d) $\frac{7}{4}$

- 31. $\tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}$ holds, when
 - (a) $x \in R$
 - (b) $x \in R (-1, 1)$ only
 - (c) $x \in R \{0\}$ only
 - (d) $x \in R [-1, 1]$ only
- 32. If $\tan A = \frac{1}{7}$, then what is $\cos 2A$ equal to?
 - (a) $\frac{24}{25}$
 - (b) $\frac{18}{25}$
 - (c) $\frac{12}{25}$
 - (d) $\frac{6}{25}$
- 33. The sides of a triangle are m, n and $\sqrt{m^2 + n^2 + mn}$. What is the sum of the acute angles of the triangle?

has both to be

医常性精神医精神 (600

- (a) 45°
- (b) 60°
- (c) 75°
- (d) 90°

- 34. What is the area of the triangle ABC with sides a=10 cm, c=4 cm and angle $B=30^{\circ}$?
 - (a) 16 cm²
 - (b) 12 cm²
 - (c) 10 cm²
 - (d) 8 cm²
- 35. Consider the following statements:
 - 1. $A = \{1, 3, 5\}$ and $B = \{2, 4, 7\}$ are equivalent sets.
 - 2. $A = \{1, 5, 9\}$ and $B = \{1, 5, 5, 9, 9\}$ are equal sets.

Which of the above statements is/are correct?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2

30 DA

- 36. Consider the following statements:
 - 1. The null set is a subset of every set.
 - 2. Every set is a subset of itself.
 - If a set has 10 elements, then its power set will have 1024 elements.

Which of the above statements are correct?

- (a) 1 and 2 only
- (b) 2 and 3 only
- (c) 1 and 3 only
- (d) 1, 2 and 3
- 37. Let R be a relation defined as xRy if and only if 2x+3y=20, where $x, y \in N$. How many elements of the form (x, y) are there in R?
 - (a) 2
 - (b) 3
 - (c) 4
 - (d) 6

- 38. Consider the following statements:
 - 1. A function $f: \mathbb{Z} \to \mathbb{Z}$, defined by f(x) = x + 1, is one-one as well as onto.
 - 2. A function $f: \mathbb{N} \to \mathbb{N}$, defined by f(x) = x + 1, is one-one but not onto.

Which of the above statements is/are correct?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2
- **39.** Consider the following in respect of a complex number Z:

1.
$$\overline{(Z^{-1})} = (\overline{Z})^{-1}$$

2.
$$ZZ^{-1} = |Z|^2$$

Which of the above is/are correct?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2

- **40.** Consider the following statements in respect of an arbitrary complex number Z:
 - 1. The difference of Z and its conjugate is an imaginary number.
 - 2. The sum of Z and its conjugate is a real number.

Which of the above statements is/are correct?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2
- **41.** What is the modulus of the complex number $i^{2n+1}(-i)^{2n-1}$, where $n \in \mathbb{N}$ and $i = \sqrt{-1}$?
 - (a) -1
 - (b) 1
 - (c) \sqrt{2}
 - (d) 2
- 42. If α and β are the roots of the equation $4x^2 + 2x 1 = 0$, then which one of the following is correct?
 - (a) $\beta = -2\alpha^2 2\alpha$
 - (b) $\beta = 4\alpha^2 3\alpha$
 - (c) $\beta = \alpha^2 3\alpha$
 - (d) $\beta = -2\alpha^2 + 2\alpha$

- **43.** If one root of $5x^2 + 26x + k = 0$ is reciprocal of the other, then what is the value of k?
 - (a) 2
 - (b) 3
 - (c) 5
 - (d) 8
- 44. In how many ways can a team of 5 players be selected from 8 players so as not to include a particular player?
 - (a) 42
 - (b) 35
 - (c) 21
 - (d) 20
- **45.** What is the coefficient of the middle term in the expansion of $(1+4x+4x^2)^5$?
 - (a) 8064
 - (b) 4032
 - (c) 2016
 - (d) 1008

46. What is $C(n, 1) + C(n, 2) + \cdots + C(n, n)$ equal to?

(a)
$$2+2^2+2^3+\cdots+2^n$$

(b)
$$1+2+2^2+2^3+\cdots+2^n$$

(c)
$$1+2+2^2+2^3+\cdots+2^{n-1}$$

(d)
$$2+2^2+2^3+\cdots+2^{n-1}$$

- **47.** What is the sum of the coefficients of first and last terms in the expansion of $(1+x)^{2n}$, where n is a natural number?
 - (a) 1
 - (b) 2
 - (c) n
 - (d) 2n
- 48. If the first term of an AP is 2 and the sum of the first five terms is equal to one-fourth of the sum of the next five terms, then what is the sum of the first ten terms?
 - (a) -500
 - (b) -250
 - (c) 500
 - (d) 250

- 49. Consider the following statements:
 - If each term of a GP is multiplied by same non-zero number, then the resulting sequence is also a GP.
 - If each term of a GP is divided by same non-zero number, then the resulting sequence is also a GP.

Which of the above statements is/are correct?

- (a) 1 only
- (b) 2 only 10 mais 13 90 fast 222
- (c) Both 1 and 2
- (d) Neither 1 nor 2
- **50.** How many 5-digit prime numbers can be formed using the digits 1, 2, 3, 4, 5 if the repetition of digits is not allowed?
 - (a) 5 was a facility with the
 - (b) 4
 - (c) 3
- **51.** If $f(x+1) = x^2 3x + 2$, then what is f(x) equal to?
 - (a) $x^2 5x + 4$
 - (b) $x^2 5x + 6$
 - (c) $x^2 + 3x + 3$
 - (d) $x^2 3x + 1$

- **52.** If x^2 , x, -8 are in AP, then which one of the following is correct?
 - (a) $x \in \{-2\}$
 - (b) $x \in \{4\}$
 - (c) $x \in \{-2, 4\}$
 - (d) $x \in \{-4, 2\}$
- **53.** The third term of a GP is 3. What is the product of its first five terms?
 - (a) 81
 - (b) 243
 - (c) 729
 - (d) Cannot be determined due to insufficient data
- 54. The element in the *i*th row and the *j*th column of a determinant of third order is equal to 2(i+j). What is the value of the determinant?
 - (a) 0
 - (b) 2
 - (c) 4

- 55. With the numbers 2, 4, 6, 8, all the possible determinants with these four different elements are constructed. What is the sum of the values of all such determinants?
 - (a) 128
 - (b) 64
 - (c) 32
 - (d) 0
- **56.** What is the radius of the circle $4x^2 + 4y^2 20x + 12y 15 = 0$?
 - (a) 14 units
 - (b) 10.5 units
 - (c) 7 units
 - (d) 3.5 units
- **57.** A parallelogram has three consecutive vertices (-3, 4), (0, -4) and (5, 2). The fourth vertex is
 - (a) (2, 10)
 - (b) (2, 9)
 - (c) (3, 9)
 - (d) (4, 10)

58. If the lines y + px = 1 and y - qx = 2 are perpendicular, then which one of the following is correct?

(a)
$$pq + 1 = 0$$

(b)
$$p+q+1=0$$

(c)
$$pq - 1 = 0$$

(d)
$$p-q+1=0$$

59. If A, B and C are in AP, then the straight line Ax + 2By + C = 0 will always pass through a fixed point. The fixed point is

$$(a)$$
 $(0, 0)$

60. If the image of the point (-4, 2) by a line mirror is (4, -2), then what is the equation of the line mirror?

(a)
$$y = x$$

$$(b) \quad y = 2x$$

(c)
$$4y = x$$

(d)
$$y = 4x$$

- **61.** Consider the following statements in respect of the points (p, p-3), (q+3, q) and (6, 3):
 - 1. The points lie on a straight line.
 - The points always lie in the first quadrant only for any value of p and q.

Which of the above statements is/are correct?

62. What is the acute angle between the lines x-2=0 and $\sqrt{3}x-y-2=0$?

63. The point of intersection of diagonals of a square *ABCD* is at the origin and one of its vertices is at *A*(4, 2). What is the equation of the diagonal *BD*?

$$(a) 2x + y = 0$$

$$(b) \quad 2x - y = 0$$

$$(c) \quad x + 2y = 0$$

$$(d) \quad x-2y=0$$

- 64. If any point on a hyperbola is $(3 \tan \theta, 2 \sec \theta)$, then what is the eccentricity of the hyperbola?
 - (a) $\frac{3}{2}$
 - (b) $\frac{5}{2}$
 - (c) $\frac{\sqrt{11}}{2}$
 - (d) $\frac{\sqrt{13}}{2}$
- 65. Consider the following with regard to eccentricity (e) of a conic section:
 - 1. e = 0 for circle
 - 2. e=1 for parabola
 - 3. e < 1 for ellipse

Which of the above are correct?

- (a) 1 and 2 only
- (b) 2 and 3 only
- (c) 1 and 3 only
- (d) 1, 2 and 3

- 66. What is the angle between the two lines having direction ratios (6, 3, 6) and (3, 3, 0)?
 - (a) $\frac{\pi}{6}$
 - (b) $\frac{\pi}{4}$
 - (c) $\frac{\pi}{3}$
 - (d) $\frac{\pi}{2}$
- 67. If l, m, n are the direction cosines of the line x-1=2(y+3)=1-z, then what is $l^4+m^4+n^4$ equal to?
 - (a) 1
 - (b) $\frac{11}{27}$
 - (c) $\frac{13}{27}$
 - (d) 4
- **68.** What is the projection of the line segment joining A(1, 7, -5) and B(-3, 4, -2) on y-axis?
 - (a) 5
 - (b) 4
 - (c) 3
 - (d) 2

69. What is the number of possible values of k for which the line joining the points (k, 1, 3) and (1, -2, k+1) also passes through the point (15, 2, -4)?

A pro 100 to 17 18

- (a) Zero
- (b) One
- (c) Two
- (d) Infinite
- 70. The foot of the perpendicular drawn from the origin to the plane x+y+z=3 is
 - (a) (0, 1, 2)
 - (b) (0, 0, 3)
 - (c) (1, 1, 1)
 - (d) (-1, 1, 3)
- 71. A vector $\vec{r} = a\hat{i} + b\hat{j}$ is equally inclined to both x and y axes. If the magnitude of the vector is 2 units, then what are the values of a and b respectively?
 - (a) $\frac{1}{2}$, $\frac{1}{2}$
 - (b) $\frac{1}{\sqrt{2}}$, $\frac{1}{\sqrt{2}}$
 - (c) $\sqrt{2}$, $\sqrt{2}$
 - (d) 2, 2

- **72.** Consider the following statements in respect of a vector $\vec{c} = \vec{a} + \vec{b}$, where $|\vec{a}| = |\vec{b}| \neq 0$:
 - 1. \vec{c} is perpendicular to $(\vec{a} \vec{b})$.
 - 2. \vec{c} is perpendicular to $(\vec{a} \times \vec{b})$.

Which of the above statements is/are correct?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2

unio de compresentar de freste a era in

- (d) Neither 1 nor 2
- 73. If \vec{a} and \vec{b} are two vectors such that $|\vec{a} + \vec{b}| = |\vec{a} \vec{b}| = 4$, then which one of the following is correct?
 - (a) \vec{a} and \vec{b} must be unit vectors.
 - (b) \vec{a} must be parallel to \vec{b} .
 - (c) \vec{a} must be perpendicular to \vec{b} .
 - (d) \vec{a} must be equal to \vec{b} .

- 74. If \vec{a} , \vec{b} and \vec{c} are coplanar, then what is $(2\vec{a} \times 3\vec{b}) \cdot 4\vec{c} + (5\vec{b} \times 3\vec{c}) \cdot 6\vec{a}$ equal to?
 - (a) 114
 - (b) 66
 - (c) 0
 - (d) -66

75. Consider the following statements:

- 1. The cross product of two unit vectors is always a unit vector.
- 2. The dot product of two unit vectors is always unity.
- The magnitude of sum of two unit vectors is always greater than the magnitude of their difference.

Which of the above statements are **not** correct?

- (a) 1 and 2 only
- (b) 2 and 3 only
- (c) 1 and 3 only
- (d) 1, 2 and 3

76. If

$$\lim_{x \to a} \frac{a^x - x^a}{x^a - a^a} = -1$$

then what is the value of a?

- (a) -1
- (b) 0
- (c) 1
- (d) 2
- 77. A particle starts from origin with a velocity (in m/s) given by the equation $\frac{dx}{dt} = x+1$. The time (in second) taken by the particle to traverse a distance of 24 m is
 - (a) ln 24
 - (b) ln 5
 - (c) 2 ln 5
 - (d) 2 ln 4

78. What is

$$\int_0^a \frac{f(a-x)}{f(x)+f(a-x)} dx$$

equal to?

- (a) a
- (b) 2a
- (c) 0
- (d) $\frac{a}{2}$

8 196 至 186 (d)。

E THE SALE SALE

79. What is

$$\lim_{x \to -1} \frac{x^3 + x^2}{x^2 + 3x + 2}$$

equal to?

- (a) 0
- (b) 1
- (c) 2
- (d) 3
- 80. If

$$\int_0^a [f(x) + f(-x)] \, dx = \int_{-a}^a g(x) \, dx$$

then what is g(x) equal to?

- (a) f(x)
- (b) f(-x) + f(x)
- (c) -f(x)
- (d) None of the above
- **81.** What is the area bounded by $y = \sqrt{16 x^2}$, $y \ge 0$ and the x-axis?
 - (a) 16π square units
 - (b) 8π square units
 - (c) 4π square units
 - (d) 2π square units

- 82. The curve $y = -x^3 + 3x^2 + 2x 27$ has the maximum slope at
 - (a) x = -1
 - (b) x = 0
 - (c) x = 1
 - (d) x=2
- 83. A 24 cm long wire is bent to form a triangle with one of the angles as 60°. What is the altitude of the triangle having the greatest possible area?
 - (a) $4\sqrt{3}$ cm
 - (b) $2\sqrt{3}$ cm
 - (c) 6 cm
 - (d) 3 cm
- **84.** If $f(x) = e^{|x|}$, then which one of the following is correct?
 - (a) f'(0) = 1
 - (b) f'(0) = -1
 - (c) f'(0) = 0
 - (d) f'(0) does not exist

85. What is

$$\int \frac{dx}{\sec x + \tan x}$$

equal to?

- (a) $\ln(\sec x) + \ln|\sec x + \tan x| + c$
- (b) $\ln(\sec x) \ln|\sec x + \tan x| + c$
- (c) $\sec x \tan x \ln |\sec x \tan x| + c$
- (d) $\ln |\sec x + \tan x| \ln |\sec x| + c$
- 86. What is

$$\int \frac{dx}{\sec^2(\tan^{-1}x)}$$

equal to?

- (a) $\sin^{-1} x + c$
- (b) $\tan^{-1} x + c$
- (c) $\sec^{-1} x + c$
- (d) $\cos^{-1} x + c$
- 87. If x+y=20 and P=xy, then what is the maximum value of P?
 - (a) 100
 - (b) 96
 - (c) 84
 - (d) 50

88. What is the derivative of

$$\sin(\ln x) + \cos(\ln x)$$

with respect to x at x = e?

(a)
$$\frac{\cos 1 - \sin 1}{e}$$

- (b) $\frac{\sin 1 \cos 1}{e}$
- (c) $\frac{\cos 1 + \sin 1}{e}$
- (d) 0
- **89.** If $x = e^t \cos t$ and $y = e^t \sin t$, then what is $\frac{dx}{dy}$ at t = 0 equal to?
 - (a) 0
 - (b) 1
 - (c) 2e
 - (d) -1
- **90.** What is the maximum value of $\sin 2x \cdot \cos 2x$?
 - (a) $\frac{1}{2}$
 - (b) 1
 - (c) 2
 - (d) 4

- **91.** What is the derivative of e^x with respect to x^e ?
 - (a) $\frac{xe^x}{ex^e}$
 - (b) $\frac{e^x}{x^e}$
 - (c) $\frac{xe^x}{x^e}$
 - (d) $\frac{e^x}{e^x}$
- **92.** If a differentiable function f(x) satisfies

$$\lim_{x \to -1} \frac{f(x)+1}{x^2-1} = -\frac{3}{2}$$

then what is $\lim_{x\to -1} f(x)$ equal to?

- (a) $-\frac{3}{2}$ for a number of equations
- (b) -1
- (c) 0
- (d) 1
- 93. If the function

$$f(x) = \begin{cases} a + bx, & x < 1 \\ 5, & x = 1 \\ b - ax, & x > 1 \end{cases}$$

is continuous, then what is the value of (a+b)?

- (a) 5
- (b) 10
- (c) 15
- (d) 20

- **94.** Consider the following statements in respect of the function $f(x) = \sin x$:
 - 1. f(x) increases in the interval $(0, \pi)$.
 - 2. f(x) decreases in the interval $\left(\frac{5\pi}{2}, 3\pi\right)$.

Which of the above statements is/are correct?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2
- **95.** What is the domain of the function $f(x) = 3^x$?
 - (a) (-∞, ∞)
 - (b) (0, ∞)
 - (c) [0, ∞)
 - (d) $(-\infty, \infty) \{0\}$
- **96.** If the general solution of a differential equation is $y^2 + 2cy cx + c^2 = 0$, where c is an arbitrary constant, then what is the order of the differential equation?
 - (a) 1
 - (b) 2
 - (c) 3
 - (d) 4

97. What is the degree of the following differential equation?

$$x = \sqrt{1 + \frac{d^2y}{dx^2}}$$

- (a) 1
- (b) 2
- (c) 3
- (d) Degree is not defined
- **98.** Which one of the following differential equations has the general solution $y = ae^x + be^{-x}$?
 - $(a) \quad \frac{d^2y}{dx^2} + y = 0$
 - $(b) \quad \frac{d^2y}{dx^2} y = 0$
 - $(c) \quad \frac{d^2y}{dx^2} + y = 1$
 - $(d) \quad \frac{dy}{dx} y = 0$
- **99.** What is the solution of the following differential equation?

$$\ln\left(\frac{dy}{dx}\right) + y = x$$

- (a) $e^x + e^y = c$
- (b) $e^{x+y} = c$
- $(c) \quad e^x e^y = c$
- $(d) e^{x-y} = c$

- 100. What is $\int e^{(2 \ln x + \ln x^2)} dx$ equal to?
 - (a) $\frac{x^4}{4} + c$
 - (b) $\frac{x^3}{3} + c$
 - (c) $\frac{2x^5}{5} + c$
 - (d) $\frac{x^5}{5} + c$
- 101. Consider the following measures of central tendency for a set of N numbers:
 - 1. Arithmetic mean
 - 2. Geometric mean

Which of the above uses/use all the data?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2
- 102. The numbers of Science, Arts and Commerce graduates working in a company are 30, 70 and 50 respectively. If these figures are represented by a pie chart, then what is the angle corresponding to Science graduates?
 - (a) 36°
 - (b) 72°
 - (c) 120°
 - (d) 168°

- 103. For a histogram based on a frequency distribution with unequal class intervals, the frequency of a class should be proportional to
 - (a) the height of the rectangle
 - (b) the area of the rectangle
 - (c) the width of the rectangle
 - (d) the perimeter of the rectangle
- 104. The coefficient of correlation is independent of
 - (a) change of scale only
 - (b) change of origin only
 - (c) both change of scale and change of origin
 - (d) neither change of scale nor change of origin
- 105. The following table gives the frequency distribution of number of peas per pea pod of 198 pods:

Number of peas	1	2	3	4	5	6	7
Frequency	4	33	76	50	26	8	1

What is the median of this distribution?

1-3-3

- (a) 3
- (b) 4
- (c) 5
- (d) 6

- 106. If M is the mean of n observations $x_1 k$, $x_2 k$, $x_3 k$, ..., $x_n k$, where k is any real number, then what is the mean of $x_1, x_2, x_3, ..., x_n$?
 - (a) M
 - (b) M+k
 - (c) M-k
 - (d) kM
- 107. What is the sum of deviations of the variate values 73, 85, 92, 105, 120 from their mean?
 - (a) -2
 - (b) -1
 - (c) 0
 - (d) 5
- **108.** Let x be the HM and y be the GM of two positive numbers m and n. If 5x = 4y, then which one of the following is correct?
 - (a) 5m = 4n
 - (b) 2m = n
 - $(c) \quad 4m = 5n$
 - (d) m = 4n

- 109. If the mean of a frequency distribution is 100 and the coefficient of variation is 45%, then what is the value of the variance?
 - (a) 2025
 - (b) 450
 - (c) 45
 - (d) 4·5
- **110.** Let two events A and B be such that P(A) = L and P(B) = M. Which one of the following is correct?
 - (a) $P(A|B) < \frac{L+M-1}{M}$
 - (b) $P(A|B) > \frac{L+M-1}{M}$
 - (c) $P(A|B) \ge \frac{L+M-1}{M}$
 - (d) $P(A|B) = \frac{L + M 1}{M}$
- 111. For which of the following sets of numbers do the mean, median and mode have the same value?
 - (a) 12, 12, 12, 12, 24
 - (b) 6, 18, 18, 18, 30
 - (c) 6, 6, 12, 30, 36
 - (d) 6, 6, 6, 12, 30

- 112. The mean of 12 observations is 75.

 If two observations are discarded, then the mean of the remaining observations is 65. What is the mean of the discarded observations?
 - (a) 250
 - (b) 125
 - (c) 120
 - (d) Cannot be determined due to insufficient data
- 113. If k is one of the roots of the equation x(x+1)+1=0, then what is its other root?
 - (a) 1
 - (b) -k
 - (c) k^2
 - $(d) -k^2$
- 114. The geometric mean of a set of observations is computed as 10. The geometric mean obtained when each observation x_i is replaced by $3x_i^4$ is
 - (a) 810
 - (b) 900
 - (c) 30000
 - (d) 81000

- 115. If $P(A \cup B) = \frac{5}{6}$, $P(A \cap B) = \frac{1}{3}$ and 118. In a binomial distribution, the mean $P(\overline{A}) = \frac{1}{2}$, then which of the following is/are correct?
 - 1. A and B are independent events.
 - A and B are mutually exclusive events.

Select the correct answer using the code given below.

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2
- 116. The average of a set of 15 observations is recorded, but later it is found that for one observation, the digit in the tens place was wrongly recorded as 8 instead of 3. After correcting the observation, the average is
 - (a) reduced by $\frac{1}{3}$
 - (b) increased by $\frac{10}{3}$
 - (c) reduced by $\frac{10}{3}$
 - (d) reduced by 50
- 117. A coin is tossed twice. If E and F denote occurrence of head on first toss and second toss respectively, then what is $P(E \cup F)$ equal to?

- (c) $\frac{3}{4}$ (d) $\frac{1}{3}$

- is $\frac{2}{3}$ and variance is $\frac{5}{9}$. What is the probability that random variable X = 2?
 - (a) $\frac{5}{36}$
 - (b)
 - (c)
- 119. If the mode of the scores 10, 12, 13, 15, 15, 13, 12, 10, x is 15, then what is the value of x?
 - (a) 10
 - (b) 12
 - (c) 13
 - (d) 15
- 120. If A and B are two events such that $P(A) = \frac{3}{4}$ and $P(B) = \frac{5}{8}$, then consider the following statements:
 - The minimum value of $P(A \cup B)$
 - The maximum value of $P(A \cap B)$

Which of the above statements is/are correct?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2

SPACE FOR ROUGH WORK

SPACE FOR ROUGH WORK

Part Hooklet Saries

TEST BOOKLET

MATHEMATICS

ot met.

A

COL : situM monthead!

Programme Annual Company of the Comp

HOSEL & D. LORDE UNA PROPERTIE ASSESSMENT OF THE MEAN PROPERTY OF THE SERVICE OF THE SERVICE ASSESSMENT OF THE SERVICE OF THE

Plants note that it is the raighbours a responsibility to encodes and till in the Lie Lead Mumber and lost Liperior Series A, it is not be a provided and without any contents a confidence plants in the OMM Antwer Steet Apy online on discrete show will center

And the state of t

the thete in prof. being a mot also to prove out on the terminosity of the control of the contro

behind the service and the service of the service o

All come time times and the second for the second f

election of the feet of the meaning of the feet of the second of the feet of t

and the conversable of the conversable of the conversable of the flower same and the examination of the flower of the flower same of the flower of the flowe

Principle for much work are depresented in the Transfer at the state and a series of the state of the series of th

a plante to measure that the property year, altered to be a measurement of the property of the

to est, in a ser service a community of the service and the service and realistic terms of the service and the

effering on the first most made large affect to the property of the design of the accordance in a principal

DO NOT GREET THE TELEFOOR TRUET RELEASE TOR OU

* * *